


**Figure 1: Transfection by injection of drops : R into G1**




A. Drop G1 on support S

➔


B. Injection of reagent R into G1

Figure 2: Transfection by fusion of drops: G1 + R



3/12

Figure 3: Transfection in drop G1 by detachment of reagent R



BEST AVAILABLE COPY

4/12

Figure 4: Fusion of cell drops G1+G2 after transfection  
*Example of the expression of a recombinant protein in a suspension of glial cells and of the activation of a suspension of neurones*





Figure 5: Photocleavage device



6/12




Figure 6





Figure 7

BEST AVAILABLE COPY

7/12

Figure 8: Membrane between two cell drops G1 and G2






FIGURE 9



9/12





FIGURE 10

BEST AVAILABLE COPY

10/12


BEST AVAILABLE COPY



**Figure 11:** Example of a spectrum obtained without CDDP and without TNF. Along the x-axis is the mass to charge ratio in Daltons (Da); along the y-axis is the signal intensity (100 corresponds to the saturation of the detector).

11/12


BEST AVAILABLE COPY



**Figure 12:** Representation of the differences between the spectra of the two phenotypes without TNF and with TNF

12/12

BEST AVAILABLE COPY



**Figure 13:** Representation of the differences between the spectra of the three phenotypes without TNF or CDDP, with TNF and with CDDP